Classical and bayesian componentwise predictors for non-compact correlated ARH(1) processes

M. Dolores Ruiz–Medina¹ and Javier Álvarez-Liébana¹

5th September 2018

¹ Department of Statistics and O. R., University of Granada, Spain. *E-mail: javialvaliebana@ugr.es*

Summary

A special class of standard Gaussian Autoregressive Hilbertian processes of order one (Gaussian ARH(1) processes), with bounded linear autocorrelation operator, which does not satisfy the usual Hilbert-Schmidt assumption, is considered. To compensate the slow decay of the diagonal coefficients of the autocorrelation operator, a faster decay velocity of the eigenvalues of the trace autocovariance operator of the innovation process is assumed. As usual, the eigenvectors of the autocovariance operator of the ARH(1) process are considered for projection, since, here, they are assumed to be known. Diagonal componentwise classical and bayesian estimation of the autocorrelation operator is studied for prediction. The asymptotic efficiency and equivalence of both estimators is proved, as well as of their associated componentwise ARH(1) plugin predictors. A simulation study is undertaken to illustrate the theoretical results derived.

Published in **REVSTAT** (in press)

https://www.ine.pt/revstat/pdf/Classicalandbayesiancomponentwise.pdf

Key words: Asymptotic efficiency; autoregressive Hilbertian processes; bayesian estimation; classical moment-based estimation; functional prediction; non-compact bounded autocorrelation operators.

1 Introduction

Functional time series theory plays a key role in the analysis of high-dimensional data (see, for example, Aue et al. [2015]; Bosq [2000]; Bosq and Blanke [2007]). Inference for stochastic processes can also be addressed from this framework (see Álvarez-Liébana et al. [2016] in relation to functional prediction of the Ornstein–Uhlenbeck process, in an ARH(1) process framework). Bosq [2000] addresses the problem of infinite–dimensional parameter estimation and prediction of ARH(1) processes, in the cases of known and unknown eigenvectors of the autocovariance operator. Alternative projection methodologies have been adopted, for example, in Antoniadis and Sapatinas [2003], in terms of wavelet bases, and Besse and Cardot [1996], in terms of spline bases. The book by Bosq and Blanke [2007] provides a general overview on statistical prediction, including Bayesian predictors, inference by projection and kernel methods, empirical density estimation, and linear processes in high–dimensional spaces (see also Blanke and Bosq [2015] on Bayesian prediction for stochastic processes). Recently, Bosq and Ruiz-Medina [2014] have derived new results on asymptotic efficiency and equivalence of classical and Bayes predictors for l^2 -valued Poisson process, where, as usual, l^2 denotes the Hilbert space of square summable sequences. Classical and Bayesian componentwise parameter estimators of the mean function and autocovariance operator, characterizing Gaussian measures in Hilbert spaces, are also compared in terms of their asymptotic efficiency, in that paper.

We first recall that the class of processes studied here could be of interest in applications, for instance, in the context of anomalous physical diffusion processes (see, for example, Gorenflo and Mainardi [2003]; Meerschaert et al. [2002]; Metzler and Klafter [2004], and the references therein). An interesting example of our framework corresponds to the case of spatial fractal diffusion operator, and regular innovations. Specifically, the class of standard Gaussian ARH(1) processes studied have a bounded linear autocorrelation operator, admitting a weak-sense diagonal spectral representation, in terms of the eigenvectors of the autocovariance operator. The sequence of diagonal coefficients, in such a spectral representation, displays an accumulation point at one. The singularity of the autocorrelation kernel is compensated by the regularity of the autocovariance kernel of the innovation process. Namely, the key assumption here is the summability of the quotient between the eigenvalues of the autocovariance operator of the innovation process and of the ARH(1) process. Under suitable conditions, the asymptotic efficiency and equivalence of the studied diagonal componentwise classical and Bayesian estimators of the autocorrelation operator are derived (see Theorem 4.1 below). Under the same setting of conditions, the asymptotic efficiency and equivalence of the corresponding classical and Bayesian ARH(1) plug-in predictors are proved as well (see Theorem 4.2 below). Although both theorems only refer to the case of known eigenvectors of the autocovariance operator, as illustrated in the simulation study undertaken in Álvarez-Liébana et al. [2017] (see also Álvarez-Liébana [2017]; Ruiz-Medina and Álvarez-Liébana [2018a]), a similar performance is obtained for the case of unknown eigenvectors, in comparison with other componentwise, kernelbased, wavelet-based penalized and nonparametric approaches adopted in the current literature (see Antoniadis and Sapatinas [2003]; Besse and Cardot [1996]; Bosq [2000]; Guillas [2001]; Mas [1999]).

Note that, for θ being the unknown parameter, in order to compute $E\{\theta|X_1,\ldots,X_n\}$, with $\{X_1,\ldots,X_n\}$ denoting the functional sample, we suppose that

$$\theta_j \bot \{ X_{i,j'}, \ i \ge 1, j' \neq j \},\$$

which leads to

$$\langle \mathbf{E} \{\theta | X_1, \dots, X_n\}, v_j \rangle_H = \mathbf{E} \{\theta_j | X_1, \dots, X_n\} = \mathbf{E} \{\theta_j | X_{1,j}, \dots, X_{n,j}\}.$$

Here, for each $j \ge 1$, $\theta_j = \langle \theta, v_j \rangle_H$, and $X_{i,j} = \langle X_i, v_j \rangle_H$, for each $i = 1, \ldots, n$, with $\langle \cdot, \cdot \rangle_H$ being the inner product in the real separable Hilbert space H. Note that $\{v_j, j \ge 1\}$ denotes an orthonormal basis of H, diagonalizing the common autocovariance operator of (X_1, \ldots, X_n) . We can then perform an independent computation of the respective posterior distributions of the projections $\{\theta_j, j \ge 1\}$, of parameter θ , with respect to the orthonormal basis $\{v_j, j \ge 1\}$ of H.

Finally, some numerical examples are considered to illustrate the results derived on asymptotic efficiency and equivalence of moment-based classical and Beta-prior-based Bayes diagonal componentwise parameter estimators, and the associated ARH(1) plug-in predictors.

2 Preliminaries

The preliminary definitions and results needed in the subsequent development are introduced in this section. We first refer to the usual class of standard ARH(1) processes introduced in Bosq [2000].

Definition 2.1 Let H be a real separable Hilbert space. A sequence $Y = \{Y_n, n \in \mathbb{Z}\}$ of H-valued random variables on a basic probability space $(\Omega, \mathcal{A}, \mathcal{P})$ is called an autoregressive Hilbertian process of order one, associated with (μ, ε, ρ) , if it is stationary and satisfies

$$X_n = Y_n - \mu = \rho(Y_{n-1} - \mu) + \varepsilon_n = \rho(X_{n-1}) + \varepsilon_n, \quad n \in \mathbb{Z},$$
(1)

where $\varepsilon = \{\varepsilon_n, n \in \mathbb{Z}\}\$ is a Hilbert-valued white noise in the strong sense (i.e., a zero-mean stationary sequence of independent H-valued random variables with $\mathbb{E}\{\|\varepsilon_n\|_H^2\} = \sigma^2 < \infty$, for every $n \in \mathbb{Z}$), and $\rho \in \mathcal{L}(H)$, with $\mathcal{L}(H)$ being the space of linear bounded operators on H. For each $n \in \mathbb{Z}$, ε_n and X_{n-1} are assumed to be uncorrelated.

If there exists a positive $j_0 \ge 1$ such that $\|\rho^{j_0}\|_{\mathcal{L}(H)} < 1$, then, the ARH(1) process in (1) is standard, and there exists a unique stationary solution to equation (1) admitting a MAH(∞) representation (see [Bosq, 2000, Theorem 3.1, p. 74]).

The autocovariance and cross-covariance operators are given, for each $n \in \mathbb{Z}$, by

$$C = E\{X_n \otimes X_n\} = E\{X_0 \otimes X_0\}, \quad D = E\{X_n \otimes X_{n+1}\} = E\{X_0 \otimes X_1\}, \quad (2)$$

where, for $f, g \in H$,

$$f \otimes g(h) = f \langle g, h \rangle_H, \quad \forall h \in H$$

defines a Hilbert–Schmidt operator on H. The operator C is assumed to be in the trace class. In particular,

$$\mathbf{E}\left\{\|X_n\|_H^2\right\} < \infty, \quad n \in \mathbb{Z}.$$

It is well-known that, from equations (1)–(2), for all $h \in H$, $D(h) = \rho C(h)$ (see, for example, Bosq [2000]). However, since C is a nuclear or trace operator, its inverse operator is an unbounded operator in H. Different methodologies have been adopted to overcome this problem in the current literature on ARH(1) processes. In particular, here, we consider the case where C(H) = H, under Assumption A2 below, since C is assumed to be strictly positive. That is, its eigenvalues are strictily positive and the kernel space of C is trivial. In addition, they are assumed to have multiplicity one. Therefore, for any $f, g \in H$, there exist $\varphi, \phi \in H$ such that $f = C(\varphi)$ and $g = C(\phi)$, and

$$\left\langle C^{-1}(f), C^{-1}(g) \right\rangle_{H} = \left\langle C^{-1}(C(\varphi)), C^{-1}(C(\phi)) \right\rangle_{H} = \left\langle \varphi, \phi \right\rangle_{H}.$$

In particular,

$$||C^{-1}(f)||_H^2 < \infty, \quad \forall f \in H.$$

Assumption A1. The operator ρ in (1) is self-adjoint with $\|\rho\|_{\mathcal{L}(H)} < 1$.

Assumption A2. The operator C is strictly positive, and its positive eigenvalues have multiplicity one. Furthermore, C and ρ admit the following diagonal spectral decompositions, such that for all $f, g \in H$,

$$C(g)(f) = \sum_{k=1}^{\infty} C_k \langle \phi_k, g \rangle_H \langle \phi_k, f \rangle_H$$
(3)

$$\rho(g)(f) = \sum_{k=1}^{\infty} \rho_k \langle \phi_k, g \rangle_H \langle \phi_k, f \rangle_H, \qquad (4)$$

where $\{C_k, k \ge 1\}$ and $\{\rho_k, k \ge 1\}$ are the respective systems of eigenvalues of C and ρ , and $\{\phi_k, k \ge 1\}$ is the common system of orthonormal eigenvectors of the autocovariance operator C.

Remark 2.1 As commented before, we consider here the case where the eigenvectors $\{\phi_k, k \ge 1\}$ of the autocovariance operator C are known. Thus, under Assumption A2, the natural way to formulate a componentwise estimator of the autocorrelation operator ρ is in terms of the respective estimators of its diagonal coefficients $\{\rho_k, k \ge 1\}$, computed from the respective projections of the observed functional data, (X_0, \ldots, X_T) , into $\{\phi_k, k \ge 1\}$. We adopt here a moment-based classical and Beta-prior-based Bayesian approach in the estimation of such coefficients $\{\rho_k, k \ge 1\}$.

From the Cauchy-Schwarz's inequality, applying the Parseval's identity,

$$\begin{aligned} \left|\rho(g)(f)\right|^2 &\leq \sum_{k=1}^{\infty} \left|\rho_k\right| \left[\langle\phi_k,g\rangle_H\right]^2 \sum_{k=1}^{\infty} \left|\rho_k\right| \left[\langle\phi_k,f\rangle_H\right]^2 \\ &\leq \sum_{k=1}^{\infty} \left[\langle\phi_k,g\rangle_H\right]^2 \sum_{k=1}^{\infty} \left[\langle\phi_k,f\rangle_H\right]^2 = \|g\|_H^2 \|f\|_H^2 < \infty. \end{aligned}$$

Thus, equation (4) holds in the weak sense.

From Assumption A2, the projection of X_n into the common eigenvector system $\{\phi_k, k \ge 1\}$ leads to the following series expansion in $\mathcal{L}^2_H(\Omega, \mathcal{A}, \mathcal{P})$:

$$X_n = \sum_{k=1}^{\infty} \sqrt{C_k} \eta_k(n) \phi_k, \quad \eta_k(n) = \frac{1}{\sqrt{C_k}} \left\langle X_n, \phi_k \right\rangle_H, \tag{5}$$

and, for each $j, p \ge 1$, and n > 0,

$$E \{\eta_j(n)\eta_p(n)\} = E \left\{ \frac{1}{\sqrt{C_j}} \langle X_n, \phi_j \rangle_H \frac{1}{\sqrt{C_p}} \langle X_n, \phi_p \rangle_H \right\}$$

$$= \frac{1}{\sqrt{C_j}} \frac{1}{\sqrt{C_p}} C(\phi_j)(\phi_p)$$

$$= \frac{1}{\sqrt{C_j}} \frac{1}{\sqrt{C_p}} C_j \langle \phi_j, \phi_p \rangle_H = \delta_{j,p},$$

where the last equality is obtained from the orthonormality of the eigenvectors $\{\phi_k, k \ge 1\}$. Hence, under **Assumptions A1–A2**, the projection of equation (1) into the elements of the common eigenvector system $\{\phi_k, k \ge 1\}$ leads to the following infinite-dimensional system of equations:

$$\sqrt{C_k}\eta_k(n) = \rho_k \sqrt{C_k}\eta_k(n-1) + \varepsilon_k(n), \quad k \ge 1,$$
(6)

or equivalently,

$$\eta_k(n) = \rho_k \eta_k(n-1) + \frac{\varepsilon_k(n)}{\sqrt{C_k}}, \quad k \ge 1,$$
(7)

where

$$\varepsilon_k(n) = \langle \varepsilon_n, \phi_k \rangle_H, \quad k \ge 1, \quad n \in \mathbb{Z}.$$

Thus, for each $j \ge 1$,

$$\{a_j(n) = \sqrt{C_j}\eta_j(n), n \in \mathbb{Z}\}$$

defines a standard AR(1) process. Its moving average representation of infinite order is given by

$$a_j(n) = \sum_{k=0}^{\infty} [\rho_j]^k \varepsilon_j(n-k), \quad n \in \mathbb{Z}.$$
(8)

Specifically, under Assumption A2,

$$E \{a_j(n)a_p(n)\} = \sum_{k=0}^{\infty} \sum_{l=0}^{\infty} [\rho_j]^k [\rho_p]^l E \{\varepsilon_j(n-k)\varepsilon_p(n-l)\}$$
$$= \sum_{k=0}^{\infty} \sum_{l=0}^{\infty} [\rho_j]^k [\rho_p]^l \delta_{k,l} \delta_{j,p} = 0, \quad j \neq p,$$
$$E \{a_j(n)a_p(n)\} = \sum_{k=0}^{\infty} \sigma_j^2 [\rho_j]^{2k}, \quad j = p,$$
(9)

where

$$\sigma_j^2 = \mathbf{E} \left\{ \varepsilon_j(n-k) \right\}^2 = \mathbf{E} \left\{ \varepsilon_j(0) \right\}^2.$$

From equation (9), under Assumptions A1–A2,

$$E\left\{ \|X(n)\|_{H}^{2} \right\} = \sum_{j=1}^{\infty} E\left\{a_{j}(n)\right\}^{2} = \sum_{j=1}^{\infty} \sigma_{j}^{2} \sum_{k=0}^{\infty} [\rho_{j}]^{2k}$$
$$= \sum_{j=1}^{\infty} \sigma_{j}^{2} \left[\frac{1}{1-[\rho_{j}]^{2}}\right] = \sum_{j=1}^{\infty} C_{j} < \infty,$$
(10)

with, as before,

$$\sum_{j=1}^{\infty} \sigma_j^2 = \mathbf{E}\left\{\|\varepsilon_n\|_H^2\right\} < \infty.$$

Equation (10) leads to the identity

$$C_j = \left[\frac{\sigma_j^2}{1 - \rho_j^2}\right], \quad j \ge 1,$$
(11)

from which, we obtain

$$\rho_k = \sqrt{1 - \frac{\sigma_k^2}{\lambda_k(C)}}, \quad \sigma_k^2 = \mathbf{E}\left\{\left\langle \phi_k, \varepsilon_n \right\rangle_H\right\}^2, \quad \forall n \in \mathbb{Z}, \quad k \ge 1.$$
(12)

Under (11), equation (7) can also be rewritten as

$$\eta_k(n) = \rho_k \eta_k(n-1) + \sqrt{1 - \rho_k^2} \frac{\varepsilon_k(n)}{\sigma_k}, \quad k \ge 1,$$

Assumption A2B. The sequences

$$\left\{\sigma_k^2, \ k \ge 1\right\}, \quad \left\{C_k, \ k \ge 1\right\}$$

satisfy

$$\frac{\sigma_k^2}{C_k} \le 1, \ k \ge 1, \quad \lim_{k \to \infty} \frac{\sigma_k^2}{C_k} = 0,$$
$$\frac{\sigma_k^2}{C_k} = \mathcal{O}(k^{-1-\gamma}), \quad \gamma > 0, \quad k \to \infty.$$
(13)

Equation (13) means that $\{\sigma_k^2, k \ge 1\}$ and $\{C_k, k \ge 1\}$ are both summable sequences, with faster decay to zero of the sequence $\{\sigma_k^2, k \ge 1\}$ than the sequence $\{C_k, k \ge 1\}$, leading, from equations (11)–(12), to the definition of $\{\rho_k^2, k \ge 1\}$ as a sequence with accumulation point at one.

Remark 2.2 Under Assumption A2B and A3 below holds.

For each $k \ge 1$, from equations (6)–(8),

$$\sum_{n=1}^{T} [\eta_k(n-1)]^2 = \frac{1}{C_k} \left[\sum_{n=1}^{T} [\varepsilon_k(n-1)]^2 + \sum_{n=1}^{T} \sum_{l=1}^{\infty} \sum_{p=1}^{\infty} [\rho_k]^l [\rho_k]^p \varepsilon_k(n-1-l) \varepsilon_k(n-1-p) \right]$$
$$= \frac{1}{C_k} \left[\sum_{n=1}^{T} [\varepsilon_k(n-1)]^2 + S(T,k) \right],$$

where

$$S(T,k) = \sum_{n=1}^{T} \sum_{l=1}^{\infty} \sum_{p=1}^{\infty} [\rho_k]^l [\rho_k]^p \varepsilon_k (n-1-l) \varepsilon_k (n-1-p)$$

Hence,
$$\sum_{n=1}^{T} [\varepsilon_k (n-1)]^2 + S(T,k) \ge 0, \text{ for every } T \ge 1, \text{ and } k \ge 1.$$

Assumption A3. There exists a sequence of real-valued independent random variables $\{\widetilde{M}(k), k \ge 1\}$ such that

$$\inf_{T \ge 1} \sqrt{\left| \frac{S(T,k)}{T\left(\sum_{n=1}^{T-1} [\varepsilon_k(n)]^2 + [\varepsilon_k(0)]^2\right)} \right|} = \inf_{T \ge 1} \sqrt{\left| \frac{\sum_{n=1}^{T} \sum_{l=1}^{\infty} \sum_{p=1}^{\infty} [\rho_k]^l [\rho_k]^p \varepsilon_k(n-1-l) \varepsilon_k(n-1-p)}{T\left(\sum_{n=1}^{T-1} [\varepsilon_k(n)]^2 + [\varepsilon_k(0)]^2\right)}} \right|} \ge [\widetilde{M}(k)]^{-1} a.s.,$$

with

$$\sum_{k=1}^{\infty} \mathbf{E}\left\{\widetilde{M}(k)\right\}^{l} < \infty, \quad 1 \le l \le 4.$$
(14)

Remark 2.3 Note that the mean value of

$$\sum_{n=1}^{T}\sum_{l=1}^{\infty}\sum_{p=1}^{\infty} [\rho_k]^l [\rho_k]^p \varepsilon_k (n-1-l)\varepsilon_k (n-1-p)$$

is of order $\frac{T\sigma_k^2}{1-(\rho_k)^2}$, and the mean value of

$$T\left(\sum_{n=1}^{T-1} \left[\varepsilon_k(n)\right]^2 + \left[\varepsilon_k(0)\right]^2\right)$$

is of order $T(T-1)\sigma_k^2$. Hence, for the almost surely boundedness of the inverse of

$$\left| \frac{S(T,k)}{T\left(\sum_{n=1}^{T-1} \left[\varepsilon_k(n)\right]^2 + \left[\varepsilon_k(0)\right]^2\right)} \right|,$$

by a suitable sequence of random variables with summable *l*-moments, for l = 1, 2, 3, 4, the eigenvalues of operator ρ must be close to one but strictly less than one. As commented in Remark 2.2, from Assumption A2B, this condition is satisfied in view of equation (12).

Assumption A4. $E \{\eta_j(m)\eta_k(n)\} = \delta_{j,k}$, with, as before, $\delta_{j,k}$ denoting the Kronecker delta function, for every $m, n \in \mathbb{Z}$, and $j, k \ge 1$.

Remark 2.4 Assumption A4 implies that the cross-covariance operator D admits a diagonal spectral decomposition in terms of the system of eigenvectors $\{\phi_k, k \ge 1\}$. Thus, under Assumption A4, the diagonal spectral decompositions (3)–(4) also hold.

The classical diagonal componentwise estimator $\widehat{\rho}_T$ of ρ considered here is given by

$$\widehat{\rho}_{T} = \sum_{k=1}^{\infty} \widehat{\rho}_{k,T} \phi_{k} \otimes \phi_{k}$$

$$\widehat{\rho}_{k,T} = \frac{\sum_{n=1}^{T} a_{k}(n-1)a_{k}(n)}{\sum_{n=1}^{T} [a_{k}(n-1)]^{2}} = \frac{\sum_{n=1}^{T} \langle X_{n-1}, \phi_{k} \rangle_{H} \langle X_{n}, \phi_{k} \rangle_{H}}{\sum_{n=1}^{T} [\langle X_{n-1}, \phi_{k} \rangle_{H}]^{2}}$$

$$= \frac{\sum_{n=1}^{T} X_{n-1,k} X_{n,k}}{\sum_{n=1}^{T} X_{n-1,k}^{2}}, \quad k \ge 1.$$
(15)

From equations (6)–(7) and (11), for each $k \ge 1$,

$$\hat{\rho}_{k,T} - \rho_{k} = \frac{\sum_{n=1}^{T} X_{n-1,k} X_{n,k}}{\sum_{n=1}^{T} [X_{n-1,k}]^{2}} - \rho_{k}$$

$$= \frac{\sum_{n=1}^{T} \rho_{k} [\eta_{k}(n-1)]^{2} + (\eta_{k}(n-1)\varepsilon_{k}(n))/\sqrt{C_{k}}}{\sum_{n=1}^{T} [\eta_{k}(n-1)]^{2}} - \rho_{k}$$

$$= \rho_{k} + \frac{\sum_{n=1}^{T} \eta_{k}(n-1)\varepsilon_{k}(n)}{\sqrt{C_{k}}\sum_{n=1}^{T} [\eta_{k}(n-1)]^{2}} - \rho_{k}$$

$$= \frac{\sum_{n=1}^{T} \eta_k(n-1)\varepsilon_k(n)}{\sqrt{\sigma_k^2/(1-\rho_k^2)} \sum_{n=1}^{T} [\eta_k(n-1)]^2}$$

$$= \sqrt{1-\rho_k^2} \frac{\sum_{n=1}^{T} \eta_k(n-1)[\varepsilon_k(n)/\sigma_k]}{\sum_{n=1}^{T} [\eta_k(n-1)]^2}.$$
 (16)

Remark 2.5 It is important to note that, for instance, unconditional bases, like wavelets, provide the spectral diagonalization of an extensive family of operators, including pseudodifferential operators, and in par-

ticular, Calderón-Zygmund operators (see Kyriazis and Petrushev [2001]; Meyer and Coifman [1997]). Therefore, the diagonal spectral representations (3)–(4), in Assumption A2, hold for a wide class of autocovariance and cross-covariance operators, for example, in terms of wavelets. When the autocovariance and the cross-covariance operators are related by a continuous function, the diagonal spectral representations (3)–(4) are also satisfied (see [Dautray and Lions, 1990, pp. 119, 126 and 140]). Assumption A2 has been considered, for example, in [Bosq, 2000, Theorem 8.5, pp. 215–216; Theorem 8.7, p. 221], to establish strong consistency, although, in this book, a different setting of conditions is assumed. Thus, Assumptions A1–A2 already have been used (e.g., in Álvarez-Liébana et al. [2017]; Bosq [2000]; Ruiz-Medina and Álvarez-Liébana [2018a]), and Assumptions A2B, A3 and A4 appear in Ruiz-Medina et al. [2016]. Assumptions A2B is needed since the usual assumption on the Hilbert-Schmidt property of ρ , made by several authors, is not considered here. At the same type, as commented before, Assumptions A2B implies Assumption A3.

The following lemmas will be used in the derivation of the main results of this paper, Theorems 4.1 and 4.2, obtained in the Gaussian ARH(1) context.

Lemma 2.1 Let $\{X_i, i = 1, ..., n\}$, be the values of a standard zero-mean autoregressive process of order one (AR(1) process) at times i = 1, ..., n, and

$$\widehat{\rho}_n = \frac{\sum_{i=1}^n \mathcal{X}_{i-1} \mathcal{X}_i}{\sum_{i=1}^n \mathcal{X}_{i-1}^2},$$

with \mathcal{X}_1 representing the random initial condition. Assume that $|\rho| < 1$, and that the innovation process is white noise. Then, as $n \to \infty$,

$$\sqrt{n} \frac{\widehat{\rho}_n - \rho}{\sqrt{1 - \rho^2}} \longrightarrow \mathcal{N}(0, 1).$$

The proof of Lemma 2.1 can be found in [Hamilton, 1994, p. 216].

Lemma 2.2 Let \mathcal{X}_1 and \mathcal{X}_2 be two normal distributed random variables having correlation $\rho_{\mathcal{X}_1\mathcal{X}_2}$, and with means μ_1 and μ_2 , and variances σ_1^2 and σ_2^2 , respectively. Then, the following identities hold:

$$E \{ \mathcal{X}_{1} \mathcal{X}_{2} \} = \mu_{1} \mu_{2} + \rho_{\mathcal{X}_{1} \mathcal{X}_{2}} \sigma_{1} \sigma_{2}$$

Var $\{ \mathcal{X}_{1} \mathcal{X}_{2} \} = \mu_{1}^{2} \sigma_{2}^{2} + \mu_{2}^{2} \sigma_{1}^{2} + \sigma_{1}^{2} \sigma_{2}^{2} + 2\rho_{\mathcal{X}_{1} \mathcal{X}_{2}} \mu_{1} \mu_{2} \sigma_{1} \sigma_{2} + \rho_{\mathcal{X}_{1} \mathcal{X}_{2}}^{2} \sigma_{1}^{2} \sigma_{2}^{2}$ (17)

(see, for example, Aroian [1947]; Ware and Lad [2003]).

Lemma 2.3 For each $k \ge 1$, the following limit is obtained:

$$\lim_{T \to \infty} T \mathbb{E} \left\{ \widehat{\rho}_{k,T} - \rho_k \right\}^2 = 1 - \rho_k^2, \quad k \ge 1$$
(18)

(see, for example, Bartlett [1946]).

3 Bayesian diagonal componentwise estimation

Now let us denote by R the functional random variable on the basic probability space $(\Omega, \mathcal{A}, \mathcal{P})$, characterized by the prior distribution for ρ . In our case, we assume that R is of the form

$$R(f)(g) = \sum_{k=1}^{\infty} R_k \langle \phi_k, f \rangle_H \langle \phi_k, g \rangle_H \ a.s., \quad \forall f, g \in H,$$

where, for $k \ge 1$, R_k is a real-valued random variable such that $R(\phi_j)(\phi_k) = \delta_{j,k}R_k$, almost surely, for every $j \ge 1$. In the following, R_k is assumed to follow a beta distribution with shape parameters $a_k > 0$ and $b_k > 0$; i.e., $R_k \sim \mathcal{B}(a_k, b_k)$, for every $k \ge 1$. We also assume that R is independent of the functional components of the innovation process $\{\varepsilon_n, n \in \mathbb{Z}\}$, and that the random variables $\{R_k, k \ge 1\}$, are globally independent. That is, for each $f, g \in H$,

$$\varphi_{R}^{f,g}(t) = E\left\{\exp\left(it\sum_{k=1}^{\infty} R_{k}\langle\phi_{k},f\rangle_{H}\langle\phi_{k},g\rangle_{H}\right)\right\}$$
$$= \prod_{k=1}^{\infty} E\left\{\exp\left(itR_{k}\langle\phi_{k},f\rangle_{H}\langle\phi_{k},g\rangle_{H}\right)\right\} = \prod_{k=1}^{\infty} \varphi_{R_{k}}\left(t\langle\phi_{k},f\rangle_{H}\langle\phi_{k},g\rangle_{H}\right).$$
(19)

Thus,

$$\varphi_R(t) = \prod_{k=1}^{\infty} \varphi_{R_k} \left(t \left(\phi_k \otimes \phi_k \right) \right),$$

where the last identity is understood in the weak-sense; i.e., in the sense of equation (19).

In the definition of R from $\{R_j, j \ge 1\}$, we can then apply the Kolmogorov extension Theorem under the condition

$$\sum_{j=1}^{\infty} \frac{a_j b_j}{(a_j + b_j + 1)(a_j + b_j)^2} < \infty$$

(see, for example, Khoshnevisan [2007]).

As in the real-valued case (see Supplementary Material 7), considering $b_j > 1$, for each $j \ge 1$, the

Bayes estimator of ρ is defined by (see Case 2 in Supplementary Material 7)

$$\widetilde{\rho}_T = \sum_{j=1}^{\infty} \widetilde{\rho}_{j,T} \phi_j \otimes \phi_j, \qquad (20)$$

with, for every $j \ge 1$,

$$\widetilde{\rho}_{j,T} = \frac{1}{2\beta_{j,T}} \left[(\alpha_{j,T} + \beta_{j,T}) \pm \sqrt{(\alpha_{j,T} - \beta_{j,T})^2 - 4\beta_{j,T}\sigma_j^2 [2 - (a_j + b_j)]} \right] \\ = \frac{\left[\sum_{i=1}^T x_{i-1,j} x_{i,j} + x_{i-1,j}^2 \right]}{2\sum_{i=1}^T x_{i-1,j}^2} \\ \pm \frac{\sqrt{\left[\sum_{i=1}^T x_{i-1,j} x_{i,j} - x_{i-1,j}^2 \right]^2 - 4\sigma_j^2 \left[\sum_{i=1}^T x_{i-1,j}^2 \right] [2 - (a_j + b_j)]}}{2\sum_{i=1}^T x_{i-1,j}^2}, \quad (21)$$

where

$$\alpha_{j,T} = \sum_{i=1}^{T} x_{i-1,j} x_{i,j}, \quad \beta_{j,T} = \sum_{i=1}^{T} x_{i-1,j}^2, \ j \ge 1, \ n \ge 2.$$
(22)

4 Asymptotic efficiency and equivalence

In this section, sufficient conditions are derived to ensure the asymptotic efficiency and equivalence of the diagonal componentwise estimators of ρ formulated in the classical (see equation (15)), and in the Bayesian (see equations (20)–(22)) frameworks.

Theorem 4.1 Under Assumptions A1-A2, A2B, A3 and A4, let us assume that the ARH(1) process X satisfies, for each $j \ge 1$, and, for every $T \ge 2$,

$$\sum_{i=1}^{T} \varepsilon_j(i) X_{i-1,j} \ge 0, \ a.s.$$
(23)

That is, $\{\varepsilon_j(i), i \ge 1\}$ and $\{X_{i-1,j}, i \ge 0\}$ are almost surely positive empirically correlated. In addition, for every $j \ge 1$, the hyper-parameters a_j and b_j of the beta prior distribution, $\mathcal{B}(a_j, b_j)$, are such that $a_j + b_j \ge 2$. Then, the following identities are obtained:

$$\lim_{T \to \infty} T \mathbf{E} \left\{ \| \widetilde{\rho}_T^- - \rho \|_{\mathcal{S}(H)}^2 \right\} = \lim_{T \to \infty} T \mathbf{E} \left\{ \| \widehat{\rho}_T - \rho \|_{\mathcal{S}(H)}^2 \right\} = \sum_{k=1}^\infty \frac{\sigma_k^2}{C_k} < \infty,$$
(24)

where $\hat{\rho}_T$ is defined in equation (15), and $\tilde{\rho}_T$ is defined from equations (20)–(22), considering

$$\widetilde{\rho}_{j,T}^{-} = \frac{1}{2\beta_{j,T}} \left[(\alpha_{j,T} + \beta_{j,T}) - \sqrt{(\alpha_{j,T} - \beta_{j,T})^2 - 4\beta_{j,T}\sigma_j^2 [2 - (a_j + b_j)]} \right],\tag{25}$$

with, as before, for each $j \ge 1$,

$$X_{i,j} = \langle X_i, \phi_j \rangle_H, \quad i = 0, \dots, T,$$

and $\alpha_{j,T}$ and $\beta_{j,T}$ are given in (22), for every $T \geq 2$.

Proof. Under Assumptions A1–A2, from Remark 8.1 and Corollary 8.1 in Supplementary Material 8, for each $j \ge 1$, and for T sufficiently large,

$$|\widehat{\rho}_{j,T}| \leq 1$$
, a.s

Also, under (23),

$$\sum_{i=1}^{T} \rho_j X_{i-1,j}^2 + \varepsilon_j(i) X_{i-1} \ge \sum_{i=1}^{T} \rho_j X_{i-1,j}^2, \quad \text{a.s.},$$

which is equivalent to

$$\widehat{\rho}_{j,T} = \frac{\sum_{i=1}^{T} \rho_j X_{i-1,j}^2 + \varepsilon_j(i) X_{i-1}}{\sum_{i=1}^{T} X_{i-1,j}^2} \ge \rho_j, \quad \text{a.s.},$$
(26)

for every $j \geq 1$.

From (26), to obtain the following a.s. inequality:

$$2|\tilde{\rho}_{j,T}^{-} - \rho_{j}| = \left| \hat{\rho}_{j,T} - \rho_{j} + 1 - \rho_{j} - \sqrt{(\hat{\rho}_{j,T} - 1)^{2} - \frac{4\sigma_{j}^{2}[2 - (a_{j} + b_{j})]}{\beta_{j,T}}} \right| \\ \leq 2|\hat{\rho}_{j,T} - \rho_{j}| \quad \text{a.s.} \quad j \ge 1,$$
(27)

it is sufficient that

$$-\widehat{\rho}_{j,T} + \rho_j \le 1 - \rho_j - \sqrt{(\widehat{\rho}_{j,T} - 1)^2 - \frac{4\sigma_j^2 [2 - (a_j + b_j)]}{\beta_{j,T}}} \le \widehat{\rho}_{j,T} - \rho_j \quad \text{a.s},$$

which is equivalent to

$$0 \le -\frac{2 - (a_j + b_j)}{\beta_{j,T}} \le 4(\hat{\rho}_{j,T} - \rho_j)(1 - \rho_j)\frac{\beta_{j,T}}{4\sigma_j^2} \quad \text{a.s.}$$
(28)

That is, keeping in mind that

$$\sigma_j^2 = C_j(1 - \rho_j^2) = C_j(1 + \rho_j)(1 - \rho_j),$$

condition (28) can also be expressed as

$$0 \le -\frac{2 - (a_j + b_j)}{\beta_{j,T}} \le 4(\hat{\rho}_{j,T} - \rho_j)(1 - \rho_j)\frac{\beta_{j,T}}{4C_j(1 + \rho_j)(1 - \rho_j)}, \quad \text{a.s.}$$

i.e.,

$$0 \le -\frac{2 - (a_j + b_j)}{\beta_{j,T}} \le (\widehat{\rho}_{j,T} - \rho_j) \frac{\beta_{j,T}}{C_j(1 + \rho_j)} \quad \text{a.s}$$

for $j \ge 1$. Since, for each $j \ge 1$,

$$\frac{\beta_{j,T}}{C_j(1+\rho_j)} \ge \frac{\beta_{j,T}}{2C_j},$$

it is sufficient that

$$0 \le -\frac{2 - (a_j + b_j)}{\beta_{j,T}} \le (\widehat{\rho}_{j,T} - \rho_j) \frac{\beta_{j,T}}{2C_j} \quad \text{a.s.}$$

$$\tag{29}$$

to hold to ensure that inequality (27) is satisfied. Furthermore, from Remark 8.1 and Corollary 8.1, in Supplementary Material 8, for each $j \ge 1$, $\beta_{j,T} \to \infty$, and

$$\beta_{j,T} = \mathcal{O}(T), \quad T \to \infty, \quad \text{a.s.}, \quad j \ge 1.$$

Also, we have, from such remark and theorem, that

_

$$(\widehat{\rho}_{j,T} - \rho_j) = \mathcal{O}(1), \quad T \to \infty, \quad \text{a.s.}, \quad j \ge 1.$$

Thus, for each $j \ge 1$, the upper bound, in (29), diverges as $T \to \infty$, which means, that, for T sufficiently large, inequality (27) holds, if $a_j + b_j \ge 2$, for each $j \ge 1$. Now, from (27), under Assumption A3, for each $j \ge 1$,

$$T|\widehat{\rho}_{j,T} - \rho_j|^2 \leq \widetilde{M}^2(j) \ a.s., \quad T|\widetilde{\rho}_{j,T} - \rho_j|^2 \leq T|\widehat{\rho}_{j,T} - \rho_j|^2 \leq \widetilde{M}^2(j) \ a.s. \tag{30}$$

Furthermore, for each $j \ge 1$, $\beta_{j,T} \to \infty$, and $\beta_{j,T} = \mathcal{O}(T)$, as $T \to \infty$, almost surely. Hence,

$$-\frac{4\sigma_j^2[2-(a_j+b_j)]}{\beta_{j,T}}\longrightarrow 0,\quad T\longrightarrow\infty,\quad \text{a.s.},\quad \forall j\ge 1.$$

From equation (25), we then have that, for each $j \ge 1$,

$$\lim_{T \to \infty} \left| \widetilde{\rho}_{j,T}^{-} - \widehat{\rho}_{j,T} \right| = \lim_{T \to \infty} \left| \frac{1}{2} \left[(\widehat{\rho}_{j,T} + 1) - \left((\widehat{\rho}_{j,T} - 1)^2 - \frac{4}{\beta_{j,T}} \sigma_j^2 [2 - (a_j + b_j)] \right)^{1/2} \right] - \widehat{\rho}_{j,T} \right| \\
= \lim_{T \to \infty} \left| \widehat{\rho}_{j,T} - \widehat{\rho}_{j,T} \right| = 0,$$
(31)

almost surely. Thus, the almost surely convergence, when $T \to \infty$, of $\tilde{\rho}_{j,T}^-$ and $\hat{\rho}_{j,T}$ to the same limit is obtained, for every $j \ge 1$.

From equation (30),

$$T[\widetilde{\rho}_{j,T} - \widehat{\rho}_{j,T}]^2 \le 2T \left[\left(\widetilde{\rho}_{j,T} - \rho_j \right)^2 + \left(\widehat{\rho}_{j,T} - \rho_j \right)^2 \right] \le 4\widetilde{M}^2(j), \quad \text{a.s.}$$
(32)

Since $\mathbb{E}\left\{\widetilde{M}^{2}(j)\right\} < \infty$, applying the Dominated Convergence Theorem, from equation (32), considering (18) we obtain, for each $j \ge 1$,

$$\lim_{T \to \infty} T \mathbb{E} \left\{ \widetilde{\rho}_{j,T} - \rho_j \right\}^2 = \lim_{T \to \infty} T \mathbb{E} \left\{ \widehat{\rho}_{j,T} - \rho_j \right\}^2 = 1 - \rho_j^2.$$
(33)

Under Assumptions A3, from (30), for each $j \ge 1$, and for every $T \ge 1$,

$$\mathbb{E}\left\{\widehat{\rho}_{j,T}-\rho_{j}\right\}^{2} \leq \mathbb{E}\left\{\widetilde{M}^{2}(j)\right\}, \quad T\mathbb{E}\left\{\widetilde{\rho}_{j,T}^{-}-\rho_{j}\right\}^{2} \leq \mathbb{E}\left\{\widetilde{M}^{2}(j)\right\}$$

with

$$\sum_{j=1}^{\infty} \mathbf{E}\left\{M^2(j)\right\} < \infty.$$

Applying again the Dominated Convergence Theorem (with integration performed with respect to a counting measure), we obtain from (33), keeping in mind relationship (12),

$$\lim_{T \to \infty} \sum_{j=1}^{\infty} T \mathbf{E} \left\{ \widetilde{\rho}_{j,T}^{-} - \rho_j \right\}^2 = \sum_{j=1}^{\infty} \lim_{T \to \infty} T \mathbf{E} \left\{ \widetilde{\rho}_{j,T}^{-} - \rho_j \right\}^2 = \sum_{j=1}^{\infty} \lim_{T \to \infty} T \mathbf{E} \left\{ \widehat{\rho}_{j,T} - \rho_j \right\}^2$$
$$= \sum_{j=1}^{\infty} 1 - \rho_j^2 = \sum_{j=1}^{\infty} \frac{\sigma_j^2}{C_j} = \lim_{T \to \infty} \sum_{j=1}^{\infty} T \mathbf{E} \left\{ \widehat{\rho}_{j,T} - \rho_j \right\}^2 < \infty,$$

in view of equation (13) in Assumption A2B. That is, equation (24) holds.

Theorem 4.2 Under the conditions of Theorem 4.1,

$$\lim_{T \to \infty} T \mathbb{E} \left\{ \| \widetilde{\rho}_T^-(X_T) - \rho(X_T) \|_H^2 \right\} = \lim_{T \to \infty} T \mathbb{E} \left\{ \| \widehat{\rho}_T(X_T) - \rho(X_T) \|_H^2 \right\} = \sum_{k=1}^{\infty} C_k (1 - \rho_k^2).$$
(34)

Here,

$$\begin{split} \widetilde{\rho}_{T}^{-}(X_{T}) &= \sum_{j=1}^{\infty} \widetilde{\rho}_{j,T}^{-} \langle X_{T}, \phi_{j} \rangle_{H} \phi_{j}, \\ \widetilde{\rho}_{\overline{j},T} &= \frac{1}{2\beta_{j,T}} \left[(\alpha_{j,T} + \beta_{j,T}) - \sqrt{(\alpha_{j,T} - \beta_{j,T})^{2} - 4\beta_{j,T}\sigma_{j}^{2}[2 - (a_{j} + b_{j})]} \right], \ j \ge 1 \\ \widehat{\rho}_{T}(X_{T}) &= \sum_{j=1}^{\infty} \widehat{\rho}_{j,T} \langle X_{T}, \phi_{j} \rangle_{H} \phi_{j}, \quad \widehat{\rho}_{j,T} \frac{\sum_{i=1}^{T} X_{i-1,j} X_{i,j}}{\sum_{i=1}^{T} X_{i-1,j}^{2}}, \quad j \ge 1 \\ \rho(X_{T}) &= \sum_{j=1}^{\infty} \rho_{j} \langle X_{T}, \phi_{j} \rangle_{H} \phi_{j}, \quad \rho_{j} = \rho(\phi_{j})(\phi_{j}), \quad j \ge 1. \end{split}$$

Proof.

From equation (31), for every $j, k \ge 1$,

$$\left[\left(\widetilde{\rho}_{j,T}^{-} - \widehat{\rho}_{j,T}\right)\left(\widetilde{\rho}_{k,T}^{-} - \widehat{\rho}_{k,T}\right)\right]^{2} \to 0, \quad \text{a.s.}, \quad T \to \infty.$$
(35)

In addition, from equation (32), for every $j, k \ge 1$,

$$\left[\left(\widetilde{\rho}_{j,T}^{-} - \widehat{\rho}_{j,T}\right)\left(\widetilde{\rho}_{k,T}^{-} - \widehat{\rho}_{k,T}\right)\right]^{2} \le 16\frac{\widetilde{M}^{2}(k)\widetilde{M}^{2}(j)}{T^{2}} \le 16\widetilde{M}^{2}(k)\widetilde{M}^{2}(j),\tag{36}$$

with

$$\mathbf{E}\left\{\widetilde{M}^{2}(k)\widetilde{M}^{2}(j)\right\} = \mathbf{E}\left\{\widetilde{M}^{2}(k)\right\}\mathbf{E}\left\{\widetilde{M}^{2}(j)\right\} < \infty,$$

under Assumption A3. Applying the Dominated Convergence Theorem from (36), the almost surely convergence in (35) implies the convergence in mean to zero, when $T \to \infty$. Furthermore, under Assumption A3, for $T \ge 2$,

$$\sum_{j=1}^{\infty} \sum_{k=1}^{\infty} T^{2} \mathbf{E} \left\{ \left(\widetilde{\rho}_{j,T}^{-} - \widehat{\rho}_{j,T} \right) \left(\widetilde{\rho}_{k,T}^{-} - \widehat{\rho}_{k,T} \right) \right]^{2} \leq 16 \left[\sum_{\substack{j,k=1\\j \neq k}}^{\infty} \mathbf{E} \left\{ \widetilde{M}^{2}(j) \right\} \mathbf{E} \left\{ \widetilde{M}^{2}(k) \right\} \right] + \left[\sum_{k=1}^{\infty} \mathbf{E} \left\{ \widetilde{M}^{4}(k) \right\} \right] < \infty.$$

$$(37)$$

From (37), for every $T \geq 2$,

$$T^{2} \mathbf{E} \left\{ \|\widetilde{\rho}_{T}^{-} - \widehat{\rho}_{T}\|_{\mathcal{S}(H)}^{4} \right\} = \sum_{j=1}^{\infty} \sum_{k=1}^{\infty} T^{2} \mathbf{E} \left\{ (\widetilde{\rho}_{j,T}^{-} - \widehat{\rho}_{j,T}) (\widetilde{\rho}_{k,T}^{-} - \widehat{\rho}_{k,T}) \right\}^{2}$$

$$\leq 16 \left[\sum_{\substack{j,k=1\\j\neq k}}^{\infty} \mathbf{E} \left\{ \widetilde{M}^{2}(j) \right\} \mathbf{E} \left\{ \widetilde{M}^{2}(k) \right\} \right]$$

$$+ \left[\sum_{k=1}^{\infty} \mathbf{E} \left\{ \widetilde{M}^{4}(k) \right\} \right] < \infty.$$
(38)

Equation (38) means that the rate of convergence to zero, as $T \to \infty$, of the functional sequence $\{\tilde{\rho}_T^- - \hat{\rho}_T, T \ge 2\}$ in the space $\mathcal{L}^4_{\mathcal{S}(H)}(\Omega, \mathcal{A}, P)$ is of order T^{-2} .

From definition of the norm in the space bounded linear operators, applying the Cauchy–Schwarz's inequality, we obtain

$$\mathbb{E}\left\{\|\widetilde{\rho}_{T}^{-}(X_{T}) - \widehat{\rho}_{T}(X_{T})\|_{H}^{2}\right\} \leq \mathbb{E}\left\{\|\widetilde{\rho}_{T}^{-} - \widehat{\rho}_{T}\|_{\mathcal{L}(H)}^{2}\|X_{T}\|_{H}^{2}\right\} \\
 \leq \sqrt{\mathbb{E}\left\{\|\widetilde{\rho}_{T}^{-} - \widehat{\rho}_{T}\|_{\mathcal{L}(H)}^{4}\right\}} \sqrt{\mathbb{E}\left\{\|X_{T}\|_{H}^{4}\right\}} \\
 \leq \sqrt{\mathbb{E}\left\{\|\widetilde{\rho}_{T}^{-} - \widehat{\rho}_{T}\|_{\mathcal{S}(H)}^{4}\right\}} \sqrt{\mathbb{E}\left\{\|X_{T}\|_{H}^{4}\right\}}.$$
(39)

From the orthogonal expansion (5) of X_T , in terms of the independent real-valued standard Gaussian random variables $\{\eta_k(T), k \ge 1\}$, we have

$$E \{ \|X_T\|_H^4 \} = \sum_{j=1}^{\infty} \sum_{k=1}^{\infty} C_j C_k E \{ \eta_j(T) \eta_k(T) \}^2 = \sum_{j=1}^{\infty} \sum_{k=1}^{\infty} C_j C_k 3 \delta_{j,k}$$

= $3 \sum_{k=1}^{\infty} C_j^2 < \infty.$ (40)

From equations (38)-(40),

$$\mathbf{E}\left\{\|\widetilde{\rho}_{T}(X_{T}) - \widehat{\rho}_{T}(X_{T})\|_{H}^{2}\right\} = \mathcal{O}\left(\frac{1}{T}\right), \quad T \to \infty.$$

Thus, $\tilde{\rho}_T^-(X_T)$ and $\hat{\rho}_T(X_T)$ have the same limit in the space $\mathcal{L}^2_H(\Omega, \mathcal{A}, \mathcal{P})$.

We now prove the approximation by Tr $(C(I - \rho^2))$ of the limit, in equation (34). Consider

$$E\left\{\|\widehat{\rho}_{T}(X_{T}) - \rho(X_{T})\|_{H}^{2}\right\} - \operatorname{Tr}\left(C(I - \rho^{2})\right) = \sum_{k=1}^{\infty} E\left\{\left(\widehat{\rho}_{k,T} - \rho_{k}\right)^{2} \eta_{k}^{2}(T)\right\} C_{k} - C_{k}(1 - \rho_{k}^{2}),$$
(41)

where

Tr
$$(C(I - \rho^2)) = \sum_{k=1}^{\infty} C_k (1 - \rho_k^2).$$

From Lemmas 2.1– 2.2 (see the last identity in equation (17)), for each $k \ge 1$, and for T sufficiently large,

$$\mathbb{E}\left\{\left(\widehat{\rho}_{k,T}-\rho_{k}\right)^{2}\eta_{k}^{2}(T)\right\} \simeq \operatorname{Var}\left\{\widehat{\rho}_{k,T}-\rho_{k}\right\}\operatorname{Var}\left\{\eta_{k}\right\}\times\left(1+2\left[\operatorname{Corr}\left(\widehat{\rho}_{k,T}-\rho_{k},\eta_{k}(T)\right)\right]^{2}\right).$$

$$(42)$$

Under Assumption A3, from equations (14)–(16), for every $k \ge 1$,

$$T\operatorname{Var}\left\{\widehat{\rho}_{k,T} - \rho_k\right\} \le \left(1 - \rho_k^2\right) \operatorname{E}\left\{\widetilde{M}^2(k)\right\}.$$
(43)

From equations (41)–(43),

$$T \mathbb{E}\left\{\|\widehat{\rho}_{T}(X_{T}) - \rho(X_{T})\|_{H}^{2}\right\} - \operatorname{Tr}\left(C(I - \rho^{2})\right) \leq \sum_{k=1}^{\infty} C_{k}(1 - \rho_{k}^{2}) \mathbb{E}\left\{\widetilde{M}^{2}(k)\right\}$$
$$\times \left[1 + 2\left[\operatorname{Corr}\left(\widehat{\rho}_{k,T} - \rho_{k}, \eta_{k}(T)\right)\right]^{2}\right]$$
$$- C_{k}(1 - \rho_{k}^{2}) \leq \sum_{k=1}^{\infty} 3C_{k} \mathbb{E}\left\{\widetilde{M}^{2}(k)\right\}$$
$$- \sum_{k=1}^{\infty} C_{k}(1 - \rho_{k}^{2}) < \infty, \qquad (44)$$

since

$$\sum_{k=1}^{\infty} C_k (1 - \rho_k^2) \le \sum_{k=1}^{\infty} C_k < \infty,$$

by the trace property of C. Here, we have applied the Cauchy–Schwarz's inequality to obtain, for a certain constant L > 0,

$$\begin{split} \sum_{k=1}^{\infty} 3C_k \mathbf{E}\left\{\widetilde{M}^2(k)\right\} &\leq 3\sqrt{\sum_{k=1}^{\infty} C_k^2 \sum_{k=1}^{\infty} \left[\mathbf{E}\left\{\widetilde{M}^2(k)\right\}\right]^2} \\ &\leq 3L\sqrt{\sum_{k=1}^{\infty} C_k \sum_{k=1}^{\infty} \mathbf{E}\left\{\widetilde{M}^2(k)\right\}} < \infty, \end{split}$$

from the trace property of C, and since

$$\sum_{k=1}^\infty {\rm E} \left\{ \widetilde{M}^2(k) \right\} < \infty,$$

under Assumption A3.

From equations (18) and (44), one can get, applying the Dominated Convergence Theorem,

$$\lim_{T \to \infty} T \mathbb{E} \left\{ \| \widehat{\rho}_T(X_T) - \rho(X_T) \|_H^2 \right\} = \sum_{k=1}^\infty C_k \lim_{T \to \infty} T \mathbb{E} \left\{ \widehat{\rho}_{k,T} - \rho_k \right\}^2$$
$$\times \lim_{T \to \infty} \left[1 + \left[\operatorname{Corr} \left(\widehat{\rho}_{k,T} - \rho_k, \eta_k(T) \right) \right]^2 \right]$$
$$= \sum_{k=1}^\infty C_k \lim_{T \to \infty} T \mathbb{E} \left\{ \widehat{\rho}_{k,T} - \rho_k \right\}^2$$
$$= \sum_{k=1}^\infty C_k (1 - \rho_k^2),$$

where we have considered that

$$\lim_{T \to \infty} \left| \operatorname{Cov} \left(\widehat{\rho}_{k,T} - \rho_k, \eta_k(T) \right) \right|^2 \leq \lim_{T \to \infty} \operatorname{E} \left\{ \widehat{\rho}_{k,T} - \rho_k \right\}^2 \operatorname{E} \left\{ \eta_k(T) \right\}^2 = \lim_{T \to \infty} \frac{1 - \rho_k^2}{T} = 0.$$

5 Numerical examples

This section illustrates the theoretical results derived on asymptotic efficiency and equivalence of the proposed classical and Bayesian diagonal componentwise estimators of the autocorrelation operator, as well as of the associated ARH(1) plug-in predictors. Under the conditions assumed in Theorem 4.1, three examples of standard zero-mean Gaussian ARH(1) processes are generated, respectively corresponding to consider different rates of convergence to zero of the eigenvalues of the autocovariance operator. The truncation order k_T in Examples 1–2 (see Sections 5.1–5.2) is fixed; i.e., it does not depend on the sample size T (see equations (46)–(47) below). While in Example 3 (see Section 5.3), k_T is selected such that

$$\lim_{T \to \infty} C_{k_T} \sqrt{T} = \infty.$$
(45)

Specifically, in the first two examples, the choice of k_T is driven looking for a compromise between the sample size and the number of parameters to be estimated. With this aim the value $k_T = 5$ is fixed, independently of T. This is the number of parameters that can be estimated in an efficient way, from most of the values of the sample size T studied. In Example 3, the truncation parameter k_T is defined as a fractional power of the sample size. Note that Example 3 corresponds to the fastest decay velocity of the eigenvalues of the autocovariance operator. Hence, the lowest truncation order for a given sample size must be selected according to the truncation rule (45). The generation of N = 1000 realizations of the functional values $\{X_t, t = 0, 1, \dots, T\}$, for

$$T = [250, 500, 750, 1000, 1250, 1500, 1750, 2000],$$

denoting as before the sample size, is performed, for each one of the ARH(1) processes, defined in the three examples below. Based on those generations, and on the sample sizes studied, the truncated empirical functional mean-square errors of the classical and Bayes diagonal componentwise parameter estimators of the autocorrelation operator ρ are computed as follows:

$$\text{EFMSE}_{\overline{\rho}_T} = \frac{1}{N} \sum_{\omega=1}^{N} \sum_{j=1}^{k_n} \left(\overline{\rho}_{j,T}^{\omega} - \rho_j \right)^2, \qquad (46)$$

$$\text{EFMSE}_{\overline{\rho}_T(X_T)} = \frac{1}{N} \sum_{\omega=1}^N \sum_{j=1}^{k_n} \left(\overline{\rho}_{j,T}^{\omega} - \rho_j \right)^2 X_{T,j}^2, \tag{47}$$

where $\overline{\rho}_{j,T}^{\omega}$ can be the classical $\hat{\rho}_{j,T}$ or the Bayes $\tilde{\rho}_{j,T}$ diagonal componentwise estimator of the autocorrelation operator, and ω denotes the sample point $\omega \in \Omega$ associated with each one of the N = 1000realizations generated of each functional value of the ARH(1) process X.

On the other hand, as assumed in the previous section,

$$\rho_k \sim \mathcal{B}(a_k, b_k), \quad a_k + b_k \ge 2, \quad a_k > 0, \quad b_k > 1,$$

for each $k \ge 1$. Thus, parameters (a_k, b_k) are defined as follows:

$$b_k = 1 + 1/100, \quad a_k = 2^k, \quad k \ge 1,$$
(48)

where

$$E\left\{\rho_k\right\} = \frac{a_k}{a_k + b_k} \to 1, \quad \operatorname{Var}\left\{\rho_k\right\} = \frac{a_k b_k}{\left(a_k + b_k + 1\right)\left(a_k + b_k\right)^2} = \mathcal{O}\left(\frac{1}{2^{2k}}\right), \quad k \to \infty,$$
(49)

with $\{\rho_k^2, k \ge 1\}$ being a random sequence such that its elements tend to be concentrated around point one, when $k \to \infty$. From (49), since

$$\sigma_k^2 = C_k \left(1 - \rho_k^2 \right), \quad k \ge 1, \tag{50}$$

Assumption A2B is satisfied. In addition, condition (23) is verified in the generations performed in the Gaussian framework.

5.1 Example 1

Let us assume that the eigenvalues of the autocovariance operator of the ARH(1) process X are given by

$$C_k = \frac{1}{k^{3/2}}, \quad k \ge 1.$$

Thus, C is a strictly positive and trace operator, where

$$\{\rho_k^2, k \ge 1\}, \{\sigma_k^2, k \ge 1\},\$$

are generated from (48)-(50).

Tables 1–2 display the values of the empirical functional mean–square errors, given in (46)–(47), associated with $\hat{\rho}_T$ and $\tilde{\rho}_T^-$, and with the corresponding ARH(1) plug–in predictors, with, as before,

$$T = [250, 500, 750, 1000, 1250, 1500, 1750, 2000],$$
(51)

considering $k_T = 5$. The respective graphical representations are displayed in Figures 1–2, where, for comparative purposes, the values of the curve 1/T are also drawn for the finite sample sizes (51).

Sample size	Classical estimator $\hat{\rho}_T$	Bayes estimator $\widetilde{\rho}_T^-$
250	$2.13(10)^{-3}$	$2.23(10)^{-3}$
500	$1.24(10)^{-3}$	$1.04(10)^{-3}$
750	$8.44(10)^{-4}$	$7.13(10)^{-4}$
1000	$6.91(10)^{-4}$	$5.84(10)^{-4}$
1250	$5.97(10)^{-4}$	$4.72(10)^{-4}$
1500	$4.89(10)^{-4}$	$3.98(10)^{-4}$
1750	$4.13(10)^{-4}$	$3.06(10)^{-4}$
2000	$3.61(10)^{-4}$	$2.59(10)^{-4}$

Table 1: Example 1. Empirical functional mean-square errors $EFMSE_{\overline{\rho}_T}$.

Sample size	Classical predictor $\hat{\rho}_T(X_T)$	Bayes predictor $\tilde{\rho}_T^-(X_T)$
250	$1.22(10)^{-3}$	$1.42(10)^{-3}$
500	$6.08(10)^{-4}$	$6.36(10)^{-4}$
750	$3.24(10)^{-4}$	$4.06(10)^{-4}$
1000	$3.05(10)^{-4}$	$2.77(10)^{-4}$
1250	$2.74(10)^{-4}$	$2.39(10)^{-4}$
1500	$2.07 (10)^{-4}$	$1.78(10)^{-4}$
1750	$1.71(10)^{-4}$	$1.48(10)^{-4}$
2000	$1.64(10)^{-4}$	$1.42(10)^{-4}$

Table 2: Example 1. Empirical functional mean-square errors $EFMSE_{\overline{\rho}_T(X_T)}$.



Figure 1: Example 1. Empirical functional mean-square estimation errors of classical (blue circle line), and Bayes (green cross line) componentwise ARH(1) parameter estimators, with $k_T = 5$, for N = 1000 replications of the ARH(1) values, against the curve 1/T (red dot line), for T = [250, 500, 750, 1000, 1250, 1500, 1750, 2000].



Figure 2: Example 1. Empirical functional mean-square prediction errors of classical (blue circle line), and Bayes (green cross line) componentwise ARH(1) plug-in predictors, with $k_T = 5$, for N = 1000 replications of the ARH(1) values, against the curve 1/T (red dot line), for T = [250, 500, 750, 1000, 1250, 1500, 1750, 2000].

5.2 Example 2

In this example, a bit slower decay velocity, than in Example 1, of the eigenvalues of the autocovariance operator of the ARH(1) process is considered. Specifically,

$$C_k = \frac{1}{k^{1+1/10}}, \quad k \ge 1.$$

Thus, C is a strictly positive self-adjoint trace operator, where $\{\rho_k^2, k \ge 1\}$ and $\{\sigma_k^2, k \ge 1\}$ are generated, as before, from (48)-(50).

Tables 3–4 show the values of the empirical functional mean–square errors, associated with $\hat{\rho}_T$ and $\tilde{\rho}_T^-$, and with the corresponding ARH(1) plug–in predictors, respectively. Figures 3–4 provide the graphical representations in comparison with the values of the curve 1/T for T given in (51), with, as before, $k_T = 5$.

Sample size	Classical estimator $\widehat{\rho}_T$	Bayes estimator $\widetilde{\rho}_T^-$
250	$4.18(10)^{-3}$	$6.09(10)^{-3}$
500	$2.20(10)^{-3}$	$2.30(10)^{-3}$
750	$1.52(10)^{-3}$	$1.39(10)^{-3}$
1000	$1.14(10)^{-3}$	$1.00(10)^{-3}$
1250	$9.55(10)^{-4}$	$7.97 (10)^{-4}$
1500	$7.97(10)^{-4}$	$6.64(10)^{-4}$
1750	$7.01(10)^{-4}$	$5.37(10)^{-4}$
2000	$6.22(10)^{-4}$	$5.00(10)^{-4}$

Table 3: Example 2. Empirical functional mean-square errors $EFMSE_{\overline{\rho}_T}$.

Table 4: Example 2. Empirical functional mean–square errors $EFMSE_{\overline{\rho}_T(X_T)}$.

Sample size	Classical predictor $\hat{\rho}_T(X_T)$	Bayes predictor $\tilde{\rho}_{T}^{-}(X_{T})$
250	$3.25(10)^{-3}$	$3.18(10)^{-4}$
500	$1.59(10)^{-3}$	$1.40(10)^{-4}$
750	$9.47(10)^{-4}$	$8.19(10)^{-4}$
1000	$7.89(10)^{-4}$	$6.88(10)^{-4}$
1250	$7.24(10)^{-4}$	$6.10(10)^{-4}$
1500	$5.53(10)^{-4}$	$4.77(10)^{-4}$
1750	$5.31(10)^{-4}$	$4.49(10)^{-4}$
2000	$4.61(10)^{-4}$	$4.00(10)^{-4}$



Figure 3: Example 2. Empirical functional mean-square estimation errors of classical (blue circle line), and Bayes (green cross line) componentwise ARH(1) parameter estimators, with $k_T = 5$, for N = 1000 replications of the ARH(1) values, against the curve 1/T (red dot line), for T = [250, 500, 750, 1000, 1250, 1500, 1750, 2000].



Figure 4: Example 2. Empirical functional mean-square prediction errors of classical (blue circle line), and Bayes (green cross line) componentwise ARH(1) plug-in predictors, with $k_T = 5$, for N = 1000 replications of the ARH(1) values, against the curve 1/T (red dot line), for T = [250, 500, 750, 1000, 1250, 1500, 1750, 2000].

5.3 Example 3

It is well-known that the singularity of the inverse of the autocovariance operator C increases, when the rate of convergence to zero of the eigenvalues of C indicates a faster decay velocity, as in this example. Specifically, here,

$$C_k = \frac{1}{k^2}, \quad k \ge 1.$$

As before, $\{\rho_k^2, k \ge 1\}$ and $\{\sigma_k^2, k \ge 1\}$ are generated from (48)-(50). The truncation order k_T satisfies

$$k_T = \lceil T^{1/\alpha} \rceil, \quad \lim_{T \to \infty} k_T = \infty, \quad \lim_{T \to \infty} \sqrt{T} C_{k_T} = \infty$$
 (52)

(see also the simulation study undertaken in Álvarez-Liébana et al. [2017], for the case of ρ being a Hilbert–Schmidt operator). In particular, (52) holds for $\frac{1}{2} - \frac{2}{\alpha} > 0$. Thus, $\alpha > 4$, and we consider $\alpha = 4.1$, i.e., $k_T = \lceil T^{1/4.1} \rceil$.

Tables 5–6 show the empirical functional mean–square errors associated with $\hat{\rho}_T$ and $\tilde{\rho}_T^-$, and with the corresponding ARH(1) plug–in predictors, respectively. As before, Figures 5–6 provide the graphical representations, and the values of the curve 1/T, for T in (51), with the aim of illustrating the rate of convergence to zero of the truncated empirical functional mean quadratic errors.

Sample size	k_T	Classical estimator $\hat{\rho}_T$	Bayes estimator $\widetilde{\rho}_T^-$
250	3	$1.73(10)^{-3}$	$1.52(10)^{-3}$
500	4	$9.72(10)^{-4}$	$1.01 (10)^{-3}$
750	5	$6.98(10)^{-4}$	$7.10(10)^{-4}$
1000	5	$5.63(10)^{-4}$	$4.35(10)^{-4}$
1250	5	$4.49(10)^{-4}$	$2.84(10)^{-4}$
1500	5	$3.94(10)^{-4}$	$2.24(10)^{-4}$
1750	6	$3.31(10)^{-4}$	$1.84(10)^{-4}$
2000	7	$3.05(10)^{-4}$	$1.70(10)^{-4}$

Table 5: Example 3. Empirical functional mean-square errors $EFMSE_{\overline{\rho}_T}$.

Table 6: Example 3. Empirical functional mean-square errors $EFMSE_{\overline{\rho}_T(X_T)}$.

Sample size	k_T	Classical predictor $\hat{\rho}_T(X_T)$	Bayes predictor $\tilde{\rho}_T^-(X_T)$
250	3	$1.92(10)^{-3}$	$1.31(10)^{-3}$
500	4	$8.24(10)^{-4}$	$5.75(10)^{-4}$
750	5	$5.60(10)^{-4}$	$4.08(10)^{-4}$
1000	5	$3.52(10)^{-4}$	$2.54(10)^{-4}$
1250	5	$2.62(10)^{-4}$	$1.45(10)^{-4}$
1500	5	$2.00(10)^{-4}$	$1.02(10)^{-4}$
1750	6	$1.37(10)^{-4}$	$9.57(10)^{-5}$
2000	6	$1.13(10)^{-4}$	$8.55(10)^{-5}$



Figure 5: Example 3. Empirical functional mean-square estimation errors of classical (blue circle line), and Bayes (green cross line) componentwise ARH(1) parameters estimators, with $k_T = \lceil T^{1/\alpha} \rceil$, $\alpha = 4.1$, for N = 1000 replications of the ARH(1) values, against the curve 1/T (red dot line), for T = [250, 500, 750, 1000, 1250, 1500, 1750, 2000].



Figure 6: Example 3. Empirical functional mean-square prediction errors of classical (blue circle line), and Bayes (green cross line) componentwise ARH(1) plug-in predictors, with $k_T = \lceil T^{1/\alpha} \rceil$, $\alpha = 4.1$, for N = 1000 replications of the ARH(1) values, against the curve 1/T (red dot line), for T = [250, 500, 750, 1000, 1250, 1500, 1750, 2000].

In Examples 1–2 in Sections 5.1–5.2, where a common fixed truncation order is considered, we can observe that the biggest values of the empirical functional mean–square errors are located at the smallest sample sizes, for which the number $k_T = 5$ of parameters to be estimated is too large, with a slightly worse performance for those sample sizes, in Example 3 in Section 5.2, where a slower decay velocity, than in Example 1, of the eigenvalues of the autocovariance operator C is considered. Note that, on the other hand, when a slower decay velocity of the eigenvalues of C is given, a larger truncation order is required to explain a given percentage of the functional variance. For the fastest rate of convergence to zero of the eigenvalues of the autocovariance operator C, in Example 3, to compensate the singularity of the inverse covariance operator C^{-1} , a suitable truncation order k_T is fitted, depending on the sample size T, obtaining a slightly better performance than in the previous cases, where a fixed truncation order is studied.

6 Final comments

This paper addresses the case where the eigenvectors of C are known, in relation to the asymptotic efficiency and equivalence of $\hat{\rho}_{j,T}$ and $\tilde{\rho}_{j,T}^-$, and the associated plug-in predictors. However, as shown in the simulation study undertaken in Álvarez-Liébana et al. [2017], a similar performance is obtained in the case where the eigenvectors of C are unknown (see also Bosq [2000] in relation to the asymptotic properties of the empirical eigenvectors of C).

In the cited references in the ARH(1) framework, the autocorrelation operator is usually assumed to belong to the Hilbert–Schmidt class. Here, in the absence of the compactness assumption (in particular, of the Hilbert–Schmidt assumption) on the autocorrelation operator ρ , singular autocorrelation kernels can be considered. As commented in the Section 1, the singularity of ρ is compensated by the regularity of the autocovariance kernel of the innovation process, as reflected in Assumption A2B.

Theorem 4.1 establishes sufficient conditions for the asymptotic efficiency and equivalence of the proposed classical and Bayes diagonal componentwise parameter estimators of ρ , as well as of the associated ARH(1) plug-in predictors (see Theorem 4.2). The simulation study illustrates the fact that the truncation order k_T should be selected according to the rate of convergence to zero of the eigenvalues of the autocovariance operator, and depending on the sample size T. Although, a fixed truncation order, independently of T, has also been tested in Examples 1–2, where a compromise between the rate of convergence to zero of the eigenvalues, and the rate of increasing of the sample sizes is found.

7 Supplementary Material: Bayesian estimation of real-valued autoregressive processes of order one

In this section, we consider the Beta-prior-based Bayesian estimation of the autocorrelation coefficient ρ in a standard AR(1) process. Namely, the generalized maximum likelihood estimator of such a parameter is computed, when a beta prior is assumed for ρ . In the ARH(1) framework, we have adopted this estimation procedure in the approximation of the diagonal coefficients { ρ_k , $k \ge 1$ } of operator ρ with respect to { $\phi_k \otimes \phi_k$, $k \ge 1$ }, in a Bayesian componentwise context. Note that we also denote by ρ the autocorrelation coefficient of an AR(1) process, since there is no place for confusion here. Let $\{X_n, n \in \mathbb{Z}\}$ be an AR(1) process satisfying

$$X_n = \rho X_{n-1} + \varepsilon_n, \quad n \in \mathbb{Z},$$

where $0 < \rho < 1$, and $\{\varepsilon_n, n \in \mathbb{Z}\}$ is a real-valued Gaussian white noise; i.e., $\varepsilon_n \sim \mathcal{N}(0, \sigma^2), n \in \mathbb{Z}$, are independent Gaussian random variables, with $\sigma > 0$. Here, we will use the conditional likelihood, and assume that (x_1, \ldots, x_n) are observed for n sufficiently large to ensure that the effect of the random initial condition is negligible. A beta distribution with shape parameters a > 0 and b > 0 is considered as a-priori distribution on ρ , i.e., $\rho \sim \mathcal{B}(a, b)$. Hence, the distribution of (x_1, \ldots, x_n, ρ) has density

$$\widetilde{L} = \frac{1}{(\sigma\sqrt{2\pi})^n} \exp\left(-\frac{1}{2\sigma^2} \sum_{i=1}^n (x_i - \rho x_{i-1})^2\right) \rho^{a-1} (1-\rho)^{b-1} \frac{\mathbf{1}_{\{0 < \rho < 1\}}}{\mathbb{B}(a,b)},$$

where

$$\mathbb{B}(a,b) = \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)}$$

is the beta function.

We first compute the solution to the equation

$$0 = \frac{\partial \ln \tilde{L}}{\partial \rho} = \frac{\partial}{\partial \rho} \left[-\frac{1}{2\sigma^2} \sum_{i=1}^n (x_i - \rho x_{i-1})^2 + (a-1) \ln \rho + (b-1) \ln(1-\rho) \right]$$

= $-\frac{1}{2\sigma^2} \sum_{i=1}^n (-2x_{i-1}(x_i - \rho x_{i-1})) + \frac{a-1}{\rho} - \frac{b-1}{1-\rho}$
= $\frac{\alpha_n}{\sigma^2} - \frac{\rho}{\sigma^2} \beta_n + \frac{a-1}{\rho} - \frac{b-1}{1-\rho},$

where

$$\alpha_n = \sum_{i=1}^n x_{i-1} x_i, \quad \beta_n = \sum_{i=1}^n x_{i-1}^2.$$

Thus, the following equation must be solved:

$$0 = \frac{\rho(1-\rho)\alpha_n}{\sigma^2} - \frac{\rho^2(1-\rho)}{\sigma^2}\beta_n + (a-1)(1-\rho) - \rho(b-1)$$

$$0 = \frac{\beta_n}{\sigma^2}\rho^3 - \frac{\alpha_n + \beta_n}{\sigma^2}\rho^2 + \left(\frac{\alpha_n}{\sigma^2} - [a+b] + 2\right)\rho + (a-1).$$

Case 1 Considering a = b = 1, and $\sigma^2 = 1$, we obtain the solution

$$\widetilde{\rho}_n = \frac{\sum_{i=1}^n x_{i-1} x_i}{\sum_{i=1}^n x_{i-1}^2}.$$

Case 2 The general case where b > 1 is more intricate, since the solutions are $\tilde{\rho}_n = 0$, and

$$\begin{split} \widetilde{\rho}_n &= \frac{1}{2\beta_n} \left[(\alpha_n + \beta_n) \pm \sqrt{(\alpha_n - \beta_n)^2 - 4\beta_n \sigma^2 [2 - (a + b)]} \right] \\ &= \frac{\sum_{i=1}^n x_{i-1} x_i + x_{i-1}^2}{2\sum_{i=1}^n x_{i-1}^2} \\ &\pm \frac{\sqrt{\left[\sum_{i=1}^n x_{i-1} x_i - x_{i-1}^2\right]^2 - 4\sigma^2 \left[\sum_{i=1}^n x_{i-1}^2\right] [2 - (a + b)]}}{2\sum_{i=1}^n x_{i-1}^2} \end{split}$$

Case 3 For $\sigma^2 = a = 1$, we have

$$\widetilde{\rho}_{n} = \frac{1}{2\beta_{n}} \left[(\alpha_{n} + \beta_{n}) \pm \sqrt{(\alpha_{n} - \beta_{n})^{2} - 4\beta_{n}(1 - b)} \right]$$

$$= \frac{1}{2\sum_{i=1}^{n} x_{i-1}^{2}} \left[\sum_{i=1}^{n} x_{i-1}x_{i} + x_{i-1}^{2} \right]$$

$$\pm \sqrt{\left[\sum_{i=1}^{n} x_{i-1}x_{i} - x_{i-1}^{2} \right]^{2} - 4\left[\sum_{i=1}^{n} x_{i-1}^{2} \right] (1 - b)}.$$

8 Supplementary Material 2: strong–ergodic AR(1) processes

This section collects some strong–ergodicity results applied in this paper, for real–valued weak– dependent random sequences. In particular, their application to the AR(1) case is considered.

A real-valued stationary process $\{Y_n, n \in \mathbb{Z}\}$ is strongly-ergodic (or ergodic in an almost surely sense), with respect to $\mathbb{E}\{f(Y_0, \dots, Y_{n-1})\}$ if, as $n \to \infty$,

$$\frac{1}{n-k}\sum_{i=0}^{n-1-k}f\left(Y_{i},\ldots,Y_{i+k}\right)\longrightarrow^{a.s.} \mathbb{E}\left\{f\left(Y_{0},\ldots,Y_{n-1}\right)\right\}, \quad k \ge 0.$$

In particular, the following lemma provides sufficient condition to get the strong–ergodicity for all second–order moments (see, for example, [Stout, 1974, Theorem 3.5.8] and [Billingsley, 1995, p. 495]).

Lemma 8.1 Let $\{\widetilde{\varepsilon}_n, n \in \mathbb{Z}\}$ be an *i.i.d.* sequence of real-valued random variables. If $f : \mathbb{R}^{\infty} \longrightarrow \mathbb{R}$ is a measurable function, then

$$Y_n = f(\widetilde{\varepsilon}_n, \widetilde{\varepsilon}_{n-1}, \ldots), \quad n \in \mathbb{Z},$$

is a stationary and strongly-ergodic process for all second-order moments.

Lemma 8.1 is now applied to the invertible AR(1) case, when the innovation process is white noise.

Remark 8.1 If $\{Y_n, n \in \mathbb{Z}\}$ is a real-valued zero-mean stationary AR(1) process

$$Y_n = \rho Y_{n-1} + \widetilde{\varepsilon}_n, \quad \rho \in \mathbb{R}, \quad |\rho| < 1, \quad n \in \mathbb{Z},$$

where $\{\widetilde{\varepsilon}_n, n \in \mathbb{Z}\}\$ is strong white noise, we can define the measurable (even continuous) function

$$f(a_0, a_1, \ldots) = \sum_{k=0}^{\infty} \rho^k a_k,$$

such that, from Lemma 8.1 and for each $n \in \mathbb{Z}$,

$$Y_n = \sum_{k=0}^{\infty} \rho^k \widetilde{\varepsilon}_{n-k} = f\left(\widetilde{\varepsilon}_n, \widetilde{\varepsilon}_{n-1}, \ldots\right),$$

is a stationary and strongly-ergodic process for all second-order moments.

In the results derived in this paper, Remark 8.1 is applied, for each $j \ge 1$, to the real-valued zeromean stationary AR(1) processes

$$\left\{X_{n,j} = \left\langle X_n, \phi_j \right\rangle_H, \ n \in \mathbb{Z}\right\},\$$

with $\{X_n, n \in \mathbb{Z}\}$ now representing an ARH(1) process.

Corollary 8.1 Under Assumptions A1–A2, for each $j \ge 1$, let us consider the real-valued zero-mean stationary AR(1) process $\{X_{n,j} = \langle X_n, \phi_j \rangle_H, n \in \mathbb{Z}\}$, such that, for each $n \in \mathbb{Z}$

$$X_{n,j} = \rho_j X_{n-1,j} + \varepsilon_{n,j}, \quad \rho_j \in \mathbb{R}, \quad |\rho_j| < 1,$$

Here, $\{\varepsilon_{n,j}, n \in \mathbb{Z}\}$ is a real-valued strong white noise, for any $j \ge 1$. Thus, for each $j \ge 1$, $\{X_{n,j}, n \in \mathbb{Z}\}$ is a stationary and strongly-ergodic process for all second-order moments. In particular, for any $j \ge 1$, as $n \to \infty$,

$$\widehat{C}_{n,j} = \frac{1}{n} \sum_{i=1}^{n} X_{i-1,j}^{2} \longrightarrow^{a.s.} \quad C_{j} = \mathbb{E} \left\{ X_{i-1,j}^{2} \right\}, \ i \ge 1,$$
$$\widehat{D}_{n,j} = \frac{1}{n-1} \sum_{i=1}^{n} X_{i-1,j} X_{i,j} \longrightarrow^{a.s.} \quad D_{j} = \mathbb{E} \left\{ X_{i-1,j} X_{i,j} \right\}, \ i \ge 1.$$

Acknowledgments

This work has been supported in part by projects MTM2012–32674 and MTM2015–71839–P (co-funded by Feder funds), of the DGI, MINECO, Spain.

References

- [Álvarez-Liébana 2017] ÁLVAREZ-LIÉBANA, J.: Functional time series: a review and comparative study. *Submitted* (2017)
- [Álvarez-Liébana et al. 2016] ÁLVAREZ-LIÉBANA, J. ; BOSQ, D. ; RUIZ-MEDINA, M. D.: Consistency of the plug-in functional predictor of the Ornstein-Uhlenbeck in Hilbert and Banach spaces. *Statist. Probab. Lett.* 117 (2016), pp. 12–22. DOI: doi.org/10.1016/j.spl.2016.04.023
- [Álvarez-Liébana et al. 2017] ÁLVAREZ-LIÉBANA, J. ; BOSQ, D. ; RUIZ-MEDINA, M. D.: Asymptotic properties of a componentwise ARH(1) plug-in predictor. J. Multivariate Anal. 155 (2017), pp. 12–34.
 DOI: doi.org/10.1016/j.jmva.2016.11.009
- [Antoniadis and Sapatinas 2003] ANTONIADIS, A. ; SAPATINAS, T.: Wavelet methods for continuoustime prediction using Hilbert-valued autoregressive processes. J. Multivariate Anal. 87 (2003), pp. 133– 158. – DOI: doi.org/10.1016/S0047-259X(03)00028-9
- [Aroian 1947] AROIAN, L. A.: The probability function of a product of two normal distributed variables. Ann. Math. Statist. 18 (1947), pp. 256–271. DOI: doi.org/10.1214/aoms/1177730442
- [Aue et al. 2015] AUE, A. ; NORINHO, D. ; HÖRMANN, S.: On the prediction of stationary functional time series. J. Amer. Statist. Assoc. 110 (2015), pp. 378–392. – DOI: doi.org/10.1080/01621459.2014.909317
- [Bartlett 1946] BARTLETT, M. S.: On the theoretical specification and sampling properties of autocorrelated time series. Supplement to J. Roy. Stat. Soc. 8 (1946), pp. 27-41. - URL http://www.jstor.org/stable/2983611
- [Besse and Cardot 1996] BESSE, P. C. ; CARDOT, H.: Approximation spline de la prévision d'un processu fonctionnel autoregréssif d'ordre 1. *Canad. J. Statist.* 24 (1996), pp. 467–487. – DOI: doi.org/10.2307/3315328
- [Billingsley 1995] BILLINGSLEY, P.: Probability and Measure, 3nd edition. John Wiley & Sons, Chicago, 1995. – ISBN 9780471007104

- [Blanke and Bosq 2015] BLANKE, D.; BOSQ, D.: Bayesian prediction for stochastic processes: Theory and applications. Sankhya A 77 (2015), pp. 79–105. DOI: doi.org/10.1007/s13171-014-0059-y
- [Bosq 2000] Bosq, D.: Linear Processes in Function Spaces. Springer, New York, 2000. ISBN 9781461211549
- [Bosq and Blanke 2007] BOSQ, D.; BLANKE, D.: Inference and predictions in large dimensions. Wiley, 2007. – ISBN 9780470017616
- [Bosq and Ruiz-Medina 2014] Bosq, D. ; RUIZ-MEDINA, M. D.: Bayesian estimation in a high dimensional parameter framework. *Electron. J. Stat.* 8 (2014), pp. 1604–1640. – DOI: doi.org/10.1214/14-EJS935
- [Dautray and Lions 1990] DAUTRAY, R.; LIONS, J.-L.: Mathematical Analysis and Numerical Methods for Science and Technology Volume 3: Spectral Theory and Applications. Springer, New York, 1990.
 - ISBN 9783642615290
- [Gorenflo and Mainardi 2003] GORENFLO, R. ; MAINARDI, F.: Fractional diffusion processes: probability distribution and continuous time random walk. *Lecture Notes in Physics* 621 (2003), pp. 148–166.
 DOI: doi.org/10.1007/3-540-44832-2_8
- [Guillas 2001] GUILLAS, S.: Rates of convergence of autocorrelation estimates for autoregressive Hilbertian processes. *Statist. Probab. Lett.* 55 (2001), pp. 281–291. – DOI: doi.org/10.1016/S0167-7152(01)00151-1
- [Hamilton 1994] HAMILTON, J. D.: Time Series Analysis. Princeton University Press, Princenton, 1994. – ISBN 9780691042893
- [Khoshnevisan 2007] KHOSHNEVISAN, D.: Probability. American Mathematical Society, UE, 2007. ISBN 9780821842157
- [Kyriazis and Petrushev 2001] KYRIAZIS, G. ; PETRUSHEV, P.: New bases for Triebel-Lizorkin and Besov spaces. Transactions of the American Mathematical Society 354 (2001), pp. 749-776. - URL http://www.ams.org/journals/tran/2002-354-02/S0002-9947-01-02916-6/S0002-9947-01-02916-6.pdf
- [Mas 1999] MAS, A.: Normalité asymptotique de l'estimateur empirique de l'opérateur d'autocorrélation d'un processus ARH(1). C. R. Acad. Sci. Paris Sér. I Math. 329 (1999), pp. 899–902.
 DOI: doi.org/10.1016/S0764-4442(00)87496-0
- [Meerschaert et al. 2002] MEERSCHAERT, M. M.; BENSON, D. A.; SCHEFFLER, H. P.; BAEUMER, B.: Stochastic solution of space-time fractional diffusion equations. *Phys. Rev. E* 65 (2002), pp. 1103–1106.
 – DOI: doi.org/10.1103/PhysRevE.65.041103

- [Metzler and Klafter 2004] METZLER, R. ; KLAFTER, J.: The restaurant at the end of the random walk: recent developments in the description of anomalous transport by fractional dynamics. J. Physics A 37 (2004), pp. R161–R208. – DOI: doi.org/10.1088/0305-4470/37/31/R01
- [Meyer and Coifman 1997] MEYER, Y.; COIFMAN, R.: Wavelets, Calderón-Zygmund and multilinear operators. Cambridge University Press, 1997. – ISBN 9780521794732
- [Ruiz-Medina and Álvarez-Liébana 2018a] RUIZ-MEDINA, M. D. ; ÁLVAREZ-LIÉBANA, J.: A note on strong-consistency of componentwise ARH(1) predictors. *Stat. Probab. Lett. (under minor revision)* (2018a)
- [Ruiz-Medina et al. 2016] RUIZ-MEDINA, M. D. ; ROMANO, E. ; FERNÁNDEZ-PASCUAL, R.: Plug-in prediction intervals for a special class of standard ARH(1) processes. J. Multivariate Anal. 146 (2016), pp. 138–150. – DOI: doi.org/10.1016/j.jmva.2015.09.001
- [Stout 1974] STOUT, W. F.: Almost sure convergence. Academic, New York-London, 1974. ISBN 9780126727500
- [Ware and Lad 2003] WARE, R. ; LAD, F.: Approximating the distribution for sums of product of normal variables. Research-Paper 2003-15. Department of Mathematics and Statistics (University of Canterbury - New Zealand) (2003). - URL http://www.math.canterbury.ac.nz/research/ucdms2003n15.pdf